

Botanical Modulation of Inflammation and Allostasis

Co-authored by
Donald R. Yance Jr., RH (AHG), CN and
Suzanne E. Sky, L.Ac., MTOM



Table of Contents

| | |
|--|-------|
| Introduction and Overview | 3 |
| Allostasis and the Inflammatory Response | 3 |
| Epigenetic Processes and Inflammation | 4 |
| Chronic Inflammation and Disease | 5 |
| Immune and Inflammatory Response | 5 |
| Inflammatory Response: Key Players | 6 |
| Natural Killer Cells, Macrophages, Cytokines | 6 |
| Tumor Necrosis Factor, NF-kB | 7 |
| COX and the Arachidonic Cascade | 8 |
| Chronic Inflammation and Epigenetic Disruption | 9 |
| Aging and Inflammation | 9 |
| Integrative Approach to Inflammation | 10 |
| Polyphenols: Powerful Phytochemicals | 11 |
| Turmeric, Green Tea, Knotweed, Resveratrol, Amla | 11-12 |
| Fatty Acids: Dietary Foundation | 12 |
| Fish Oils, Sea Buckthorn Oil | 13 |
| Herbs and Natural Compounds to Calm Inflammation and Pain | 13 |
| Frankincense, Feverfew, Magnolia, Andrographis, Ginger, Peppers/Piperine | 14-15 |
| Corydalis, Willow Bark, Notoginseng, White Peony, Dong Quai | 15-17 |
| Conclusion | 17 |
| References | 18-25 |
| Author Bios | 26 |

Introduction and Overview


Allostasis and the Inflammatory Response

Inflammation is an intrinsically protective response of the immune system to injury, infection or invasion by microorganisms that usually leads to resolution and tissue healing within a short period of time. The immune response involves highly complex interactions between many signaling networks, molecules and systems that modulate cellular homeostasis and health while responding to systemic challenges.¹⁻⁴

Our immune system is comprised of a vast network of cells and tissues connected through ongoing intercellular communication with intricate cascades of biochemical transformations mediated by enzymes and many other factors. The study of immune and inflammatory response spans multiple scientific disciplines including cell biology, biochemistry, biology and genetics. Current research explores the intricate interrelationship between inflammatory, epigenetic and allostatic processes. Continual communication and feedback loops facilitate the immune, cardiovascular, endocrine, epigenetic and neurological systems to work in an integrated manner to maintain stability. This process of “achieving stability through change” is the constant adaptive process of allostasis: the ongoing, active process by which the body maintains homeostasis while responding to ongoing events and challenges.¹ The epigenetic, immune and inflammatory responses are but a few aspects of this ongoing process of allostasis.

Complex inflammatory processes occur through multiple mechanisms and mediators leading to vasodilation, increased vascular permeability, migration of white blood cells to the affected area and many other events.⁵ Acute inflammation is usually resolved within a short period of time - often around two weeks. When the inflammatory response becomes abnormally prolonged this creates wear and tear on the cells, tissues and organs of the body, which creates what is referred to as allostatic overload and eventually resulting in disease.¹ What kind of disease and where it occurs depends on which cellular systems are damaged due to the ongoing process of chronic inflammation.

An example of this complex interrelationship is seen with cytokines. There are both pro- and anti-inflammatory cytokines, which are produced by many cells throughout the body. Cytokines influence each other and are also regulated by glucocorticoids and catecholamines, both of which can increase production of pro-inflammatory cytokines. Glucocorticoids can also inhibit this production and their influence is found to depend on factors such as dosage along with cell or tissue type.



Nonlinear network of mediators of allostasis involved in the stress response. Arrows show the multiple pathways for regulation and indicate that each system regulates the others in a reciprocal manner, creating a non-linear network.

McEwen BS. Protective and damaging effects of stress mediators: central role of the brain. Dialogues in Clinical Neuroscience 2006. Vol 8(4):367-381.

The balance between parasympathetic and sympathetic nervous system engagement plays a role in the regulatory process of inflammation. Generally, the parasympathetic nervous system tends to engender


healing and repair and is found to exert an anti-inflammatory influence. Scientists find that the mediators of these systems function in a nonlinear manner. This means that when there is a change (increase or decrease) in one mediator, all the other mediators respond in a compensatory manner. These changes are dependent on multiple factors such as time and level of change of the mediators.¹

Multiple stress factors, especially over time, can lead to allostatic overload. For example, loss of sleep causes multiple physiological changes including lessened parasympathetic tone, increased evening cortisol and insulin levels along with an increase in proinflammatory cytokine levels and oxidative stress markers.¹

Due to the degree of complexity and continual interaction of intercellular communication throughout physiological systems, only a few of the multitudinous components and interactions involved with the inflammatory response will be discussed here.

Epigenetic Processes and Inflammation

Epigenetics explores the role of exogenous and endogenous factors on gene expression. Our health is not determined by our genes per se but rather by their response to stimuli. As one of the basic cellular regulation systems, epigenetic control plays key roles in regulating inflammatory and immune responses.³ The combined influence of transcription factors (by regulatory gene proteins) and epigenetic mechanisms governs transcriptional control of gene expression. For example, in a study of the inflammatory processes of atherosclerosis, it is found that epigenetic modulators regulate vascular, immune and tissue-specific gene expression within the atherosclerotic lesion. They are also involved in transcriptional regulation of immune cells in the vascular wall, which modulates inflammation through expression of cytokines and chemokines.³



Schematic representation of plaque development.

Wierda RJ, Geutskens SB, et al. *Epigenetics in atherosclerosis and inflammation.* J Cell Mol Med. Epigenetics Review Series 2010. Vol 14(6A):1225-1240

Epigenetic processes are key factors in regulation of gene expression and cytokine expression.³ Epigenetic modifications are mediated by various enzymes responsible for maintaining the epigenome (the overall epigenetic state of an organism), which is critical to normal development and health.⁶ Multiple components influence epigenetic regulation of genes and cause modification of genes. Since epigenetic alterations accumulate over time, environmental, dietary and other factors exert a powerful influence on genetic

expression. Cell differentiation, such as monocyte differentiation into macrophages in the immune response and cell-specific gene expression, is determined in large part through epigenetic mechanisms.³

Scientists are interested in epigenetic processes because they are reversible and seen as a promising therapeutic target due to their inherent plasticity.^{3, 6-8} Botanicals are of particular interest in this regard. There is a large body of research discussing the link between inflammation and reprogramming of the epigenome.^{3,4; 6-8} For example, curcumin is found to act as an inhibitor of an epigenetic-mediating enzyme and through this mechanism to significantly influence the favorable LDL/HDL ratio.³

Chronic Inflammation and Disease

Decades of intensive research find a direct relationship between chronic inflammation, aging processes and disease. A persistent, dysfunctional inflammatory response that continues over time damages cells, disrupts cell-signaling networks, causes tissue damage and can predispose a person to autoimmune conditions or other diseases. Chronic inflammation is implicated in the pathogenesis of many conditions including atherosclerosis, arthritis, rheumatoid arthritis, chronic obstructive pulmonary disease (COPD), cancer, stroke, chronic pain, periodontal disease, neurodegenerative conditions and cardiovascular disease. Ongoing research identifies and maps the specific players, processes and targets in the inflammatory response and aims to discover compounds that can restore allostasis.⁹⁻¹¹

Increased oxidative and genotoxic stress contribute to a vicious cycle of inflammation and oxidative stress that provokes cellular damage, epigenetic disturbances and a cellular environment where senescent cells create molecules that cause further disturbances for surrounding cells. Studies show that DNA damage in aging cells can activate the innate immune system release of inflammatory cytokines and induce activation of the NF-kB (nuclear factor-kappa beta) system. This chronic inflammatory response increases oxidative stress, lipid peroxidation and induces secretion of other deleterious products. This in turn causes degeneration of the cellular matrix and can trigger tissue-specific age-related degenerative conditions.² Inflammatory processes involving NF-kB, tumor necrosis factor (TNF) and cytokines are found to play critical roles in the development of inflammatory conditions including atherosclerosis, rheumatoid arthritis, irritable bowel disease, asthma and COPD.¹²

Immune and Inflammatory Responses

The two aspects of the immune system – innate and adaptive – function together in a highly integrated manner. Innate defenses respond directly to challenges while the adaptive defense system develops its response after initial encounter with a challenging molecule. The innate system, the body's first responder, instantly recognizes microbial, viral or other pathogens through mechanisms known as pattern recognition receptors. Toll proteins (toll-like receptors) are key in recognizing foreign microbes and activating innate cellular response.¹³ The innate immune system calls the adaptive system into play. Adaptive system responses, highly specific to the pathogen that induced them, act through antibody and cell-mediated responses.^{13,14}

The acute phase of inflammation is characterized by an influx of white blood cell components. These help neutralize and eliminate any invading microorganisms, help prevent infection and begin the process of healing and repair. The four cardinal signs of inflammation are redness, heat, swelling and pain. Loss of function can also occur. White blood cell components including macrophages, lymphocytes and leukocytes increase in numbers to play a primary role in the inflammatory process. During the resolution phase, these molecules return to normal numbers and function. Acute inflammation usually resolves within about two weeks.¹³



Some of the key compounds that promote the inflammatory process also play a role in the resolution of inflammation, helping restore healthy function and homeostasis.^{5,9-11,15} These include enzymes (such as cyclooxygenase), lipids (prostaglandins) and proteins (cytokines and others).

Inflammatory Response: Key Players

Natural Killer (NK) Cells

NK cells are a form of lymphocytes produced in and released from the bone marrow. They are a key player in the innate immune response and are mainly found in the blood circulation and in the spleen. In addition, they are cytotoxic and are able to directly kill problematic cells. NK cells modulate expression of cytokine and chemokine receptors.¹⁶

Macrophages

Innate defenses are mediated by NK cells and by phagocytic cells such as neutrophils and macrophages (MPs). MPs, which are found throughout the body and named according to their location. For example, Kupffer cells are found in the liver and microglial cells are found in the brain. Microglial cells play a major role in modulating both pain and inflammation.¹⁷⁻¹⁹

MPs, often the first responders of the immune system, perform a variety of functions. They can act as phagocytes to eliminate apoptotic cells, digest waste products from tissues, and help promote wound healing. MPs are essential for immunity, overall development and tissue homeostasis.²⁰

Reactive oxidative species (ROS), produced by MPs, damage an invading microbe's cell wall, which makes it more permeable to attack and able to be neutralized and removed from the body. Hence, ROS formation and subsequent oxidative stress is linked with the inflammatory process.

During the inflammatory response, MPs secrete cytokines along with tumor necrosis factor. Cytokines activate white blood cells, including lymphocytes and neutrophils, and promote inflammation. The release and activity of cytokines is modulated by cell-signaling and through highly complex intercellular processes involving specialized membranes and molecules. When this inflammatory response becomes chronic or excessive due to dysregulation, it disrupts cellular function and contributes to disease formation.^{13;20-22}

Cytokines

MPs communicate information to other immune cells through signaling molecules known as cytokines, which are secreted by MPs and by other immune system cells. Cytokines are important mediators of both the innate and specific defense systems. Often regarded as the hormones of the immune system, their role is considered as vital to life as that of hormones and neurotransmitters.^{20,23}

Cytokines comprise a huge family of molecules that are classified according to their source and function. Cytokines provide molecular cues and messaging that help maintain physiological stability and communication between organ systems. Specific cytokines stimulate stem-cell growth, proliferation and cell differentiation. They can function as growth factors and growth inhibitors.^{13,20,24} Some cytokines function as epigenetic regulators.^{4,20} Binding to cell-surface receptors, they generate a cell-signaling cascade that

influences cell function, along with regulation of certain genes and their transcription factors.²⁰

Cytokines are found to play a key role in signaling the brain; inducing it to produce neuro-chemical, neuro-endocrine, neuro-immune and behavioral changes as part of the immune response.²³ Many types of cytokines are synthesized and released within the central nervous system and in the brain where they are secreted by astrocytes and microglia. In some cases, neurons can also produce cytokines.

Cytokines modulate the HPAT (hypothalamic-pituitary-adrenal-thyroid) axis. Many symptoms that occur during inflammation and infection are attributed in part to the influence of cytokine activity including increased sleep and decreased appetite. Thus cytokines can act as immune-modulators and neuro-modulators, influencing metabolic and other pathways to maintain homeostasis and restore health.²³

Chemokines and interleukins (ILs) are two interrelated members of the vast cytokine family. Chemokines are unique cytokines released by many different kinds of cells, most often induced by pro-inflammatory cytokines including TNF and IL-6.²⁰ ILs are a large, diverse group of compounds with implications too complex to discuss here. Generally, they are immune-modulators. IL-6 plays a role in both pro-and anti-inflammatory processes in functions including immunity, tissue repair and metabolism.²⁰


Tumor Necrosis Factor (TNF)

TNF (formerly known as TNF- α) is a glycoprotein that stimulates the acute phase of the immune response and regulates chemokine release. In the hypothalamus, TNF stimulates the release of corticotropin releasing hormone. A key factor in elevating synthesis of C-reactive protein and other mediators, it exerts many more effects as a major inducer of the inflammatory cascade. Excess TNF is found to play a role in conditions as diverse as irritable bowel syndrome, psoriasis, rheumatoid arthritis, asthma, infectious diseases, cancer and a number of autoimmune conditions.²⁰

NF- κ B (nuclear factor kappa-beta)

NF- κ B, a major driver in the inflammatory process, is a transcription protein. Researchers consider it to be an ancient signaling pathway system that plays a key role in the immune system with a vast influence on health and aging.² Activation mechanisms and responses are unique and specific for different cell types. The NF- κ B system is a pleiotropic regulator that has both beneficial and detrimental responses. One of its beneficial functions lies in its key role in the innate immune response.

The NF- κ B signaling system is considered a focal point linking the body's immune response and its transcriptionally-mediated adaptive response. NF- κ B is a transcription factor and a mediator of gene expression. It is considered to be a cytoplasmic sensor that is activated by immune attacks and by many other signals including oxidative or genotoxic stress and tissue injury.² Through various mechanisms, NF- κ B plays a role in controlling the magnitude and duration of the inflammatory response.¹² NF- κ B regulates gene



Cytokine regulation in health and disease.

Kronfol Z, Remick DG. *Cytokines and the brain: implications for clinical psychiatry.* Am J Psychiatry 2000. 157:683-694.


expression and influences cell health.¹⁰ Prolonged or excessive activation is linked with formation of disease processes.² NF-κB overexpression is linked to expression of multiple disorders including arthritis and cellular disturbances.^{25,26} As a transcription factor, NF-κB can stimulate growth genes, promoting uncontrolled cellular growth.²⁷

COX (cyclooxygenase) and the Arachidonic Acid (AA) Cascade

The COX-1 and COX-2 pathways are major players in the inflammatory cascade. COX is the key enzyme required to convert AA (arachidonic acid) to PGs (prostaglandins). While COX-1 is implicated in the inflammatory process it also plays a role in homeostasis and health, particularly in the kidneys and gastric mucosa. COX-2 induces the inflammatory cascade, but is sometimes involved in resolution of inflammation.^{15,28} COX-2 is found to modulate brain health, neural response, development, imprinting and adaptation. However, chronic activation of COX-2 provokes numerous disease processes.¹⁵


AA is transformed by COX-2 enzymes to PGs and thromboxane. PGs along with the compounds thromboxanes, leukotrienes and lipoxins are collectively referred to as eicosanoids. Eicosanoids are significantly elevated during the inflammatory response.^{29,30}

PGs are phospholipid compounds that act as biological regulators; they respond contextually to either promote inflammation or enhance resolution of inflammation. They trigger signal transduction pathways, influence immune response and increase sensitivity of pain receptors.¹³ PGs can support physiological homeostasis through modulating intracellular cell-signaling pathways.⁸ PGs mediate inflammatory processes including cytokine production.^{10,12,31}



NF-κB system is at the hub of the aging network.

Salminen A, Kaarniranta K. Genetics vs. entropy: Longevity factors suppress the NF-κB-driven entropic aging process. *Aging Research Reviews* 2010. (9)298-314.



Products and enzymes of arachidonic acid metabolism involved in the inflammatory process.

Martel-Pelletier J, Lajeunesse D et al. Therapeutic role of dual inhibitors of 5-LOX and COX, selective and non-selective non-steroidal anti-inflammatory drugs. *Ann Rheum Dis* 2003. 762: 501-509.

Chronic Inflammation and Epigenetic Disruption

Factors that contribute to chronic inflammatory processes and cellular dysfunction include stress (chemical, physical and psychological), viral infection, bacterial imbalance (i.e. *Helicobacter pylori*), environmental pollutants (air pollution, smoke, etc.) and dietary contaminants (everything from fried foods to GMOs, xenobiotics and hormone/pesticide contamination). These incite the expression of inflammatory processes including reactive oxidative species, TNF, IL factors, NF-kB, COX-2 and 5-LOX, matrix metalloproteinases and chemokines. Chronic inflammatory conditions correlate with multiple adverse epigenetic changes including changes in genomic markers and cellular transcription dysfunctions.^{4,6}

Epigenetic mechanisms modulate the expression of many pro-inflammatory factors including TNF- α , interleukins, tumor suppressor genes and oncogenes. They are found to modulate autocrine and paracrine activation of NF-kB. The inflammatory response is interwoven with the body's regulatory network involving signal- and gene-specific levels. Specific genes are activated for immune response, antimicrobial defense, and for tissue repair and remodeling.

Inflammatory genes are regulated by a number of transcription factors including NF-kB and by epigenetic activities such as DNA methylation.⁶ Overexpression of transcription factors such as NF-kB is one of the key drivers responsible for expression of these genes. Over time, tissue changes and epigenetic disruption can occur at the sites of chronic inflammation.⁴ For example, rheumatoid arthritis (RA) is an inflammatory condition with progressive degeneration of articular cartilage and bone. Epigenetic factors are found to contribute to RA.⁶ Research findings estimate that up to 25% of all cancers are linked with chronic inflammation and its manifestations of cellular dysfunction including transcription errors, aberrant epigenetic processes (which influence normal gene expression), DNA methylation and other processes.⁴

Aging and Inflammation

The aging process has long been linked with inflammation and specifically with pro-inflammatory changes in cellular status. Multiple changes are noted in both the innate and adaptive immune response systems.¹⁶ The combination of inflammation, decline in adaptive immunity, and changes in cell function and cell-signaling are thought to facilitate unhealthy aging and promote formation of degenerative diseases.²

Researchers report that with aging cytokines are often elevated and that IL-6 and TNF are upregulated. TNF regulates adaptive immunity and contributes to its decline with aging (immunosenescence). In addition, studies show that insulin and insulin-like growth factor signaling can stimulate the NF-kB system, which enhances the inflammatory response and exerts other negative influences on the aging process.² Research finds that the NF-kB system plays a key role in mediating aging and is overactive in many age-related, chronic inflammatory diseases.^{2,12}

Members of the TNF family regulate many factors including age-related muscle atrophy (sarcopenia) and the cachexia that is seen in inflammatory conditions.² In both these conditions, the TNF signaling pathway is mediated through the activation of the NF-kB system. Multiple signaling pathways, endocrine regulation and metabolic factors are also implicated in muscle atrophy. Other members of the TNF family play a role in the catabolic aging process.² Inflammation is linked to frailty syndrome in the aging, which is characterized as dysregulation in multiple systems and low physiological reserves. Upregulated expression of stress-responsive genes in the inflammatory pathways is found to contribute to frailty syndrome.^{16,32,33}


Integrative Approach to Inflammation

Multiple studies demonstrate the effectiveness of stress reduction, mindfulness, meditation practices and a healthy diet to mediate inflammation. Nutrigenomics integrates the science of genomics with that of nutrition to study the influence of dietary components on genes.³⁴

Plant-based foods, herbs and spices contain multiple vitamins, minerals and phytonutrients essential to human health. Most of these plants have been used for thousands of years as part of the human diet and as medicines. Modern scientists are avidly studying these plants to discover their bioactive compounds in search for new answers to healing inflammation and helping to restore and maintain health.

Some of the compounds of greatest interest include the polyphenols, phytoestrogens, saponins, terpenoids, isothiocyanates, phytosterols, carotenoids and omega-3 fatty acids.³⁵ Phytonutrients such as these and others are found to modulate healthy inflammatory processes and to help restore normal cell function and cellular homeostasis. They are able to calm oxidative stress and inflammatory pathways including the NF-kB, LOX and COX pathways.³⁶

Traditionally, researchers and clinicians have focused on targeting specific pathways, such as the well-known COX-2, in order to address inflammatory conditions. Currently, scientists are exploring targeting multiple pathways. Botanicals are of great interest due to their natural multi-target ability. For example, herbs such as Turmeric are found to exert far-reaching influence on multiple pathways. Curcumin can influence and modulate cytokines, enzymes, transcriptional factors, receptors and multiple other factors.³⁷



This diagram, while it focuses on curcumin, illustrates the fact that plant polyphenolic compounds work through multiple pathways to modulate cellular health and function.

Anand P, Sundaram C, et al. Curcumin and cancer: An "old-age" disease with an "age-old" solution. Cancer Letters 267 (2008) 133–164.

Botanical medicines contain a wide array of natural compounds beneficial to mediate chronic inflammation and epigenetics to support the return of dysregulated biochemical pathways to homeostasis and health. Many botanicals are found to successfully calm NF-kB and to modulate cytokine expression.^{2,10,12,24}

Polyphenols: Powerful Phytochemicals

Botanical polyphenols are found to alleviate inflammatory processes and to protect against age-related degeneration. Many polyphenols inhibit NF-kB activation and are potent antioxidants.^{2,31} Polyphenols are a vast family of naturally-occurring phytochemicals produced by plants as part of their defense system. Almost half of the 10,000 known polyphenolic compounds are flavonoids. Flavonoids are classified into six major subgroups: flavones, flavonols, isoflavones, flavones, anthocyanidins and flavanols (catechins and proanthocyanadins).³⁸

Polyphenols are abundant in vegetables, fruits, whole grains, herbs and spices. Epidemiological studies demonstrate an inverse relationship between a polyphenolic-rich diet and risk of chronic human diseases including cardiovascular, diabetes, osteoporosis, chronic inflammation, cancer and neurodegenerative conditions.^{3,9,39} Plant phenolic compounds are widely studied for their role in preventing both occurrence and progression of tumor growth and are found to inhibit mutagenesis and carcinogenesis in humans.^{40,41} They are also found to be cardio-protective, neuro-protective and cyto-protective.

Plant polyphenols protect cellular health through many actions such as scavenging free radicals, antioxidant and anti-inflammatory activity. They exert a powerful influence to normalize cell-signaling and to support healthy cellular function^{40,41} including apoptosis (normal cell death), cellular metabolism and maintenance of healthy cellular structure.³⁹ Plant polyphenols modulate cell-signaling pathways and demonstrate the ability to modulate enzyme response to help decrease oxidative stress.^{9,10,12,31,39,40}

Turmeric (*Curcuma longa*) and Wild Turmeric (*Curcuma aromatica*)

Turmeric rhizome, one of the most profoundly researched herbs, is found to work through multiple pathways. For centuries it has been world-renowned as a cooking spice, food, herbal medicine and dye. Traditional medicines prize these herbs for multiple benefits and they are noted for their anti-inflammatory influence, which is validated by modern research.⁴²

Turmeric's active constituents, curcuminoids, are yellowish-orange, lipid-soluble natural phenolic compounds. They demonstrate many protective health benefits, offering powerful anti-inflammatory, antioxidant, antineoplastic, antiviral and immune-modulating activity.¹⁰ Curcumin works to calm inflammatory pathways including NF-kB, TNF and IL factors^{43,44} and inhibits COX-2 and 5-LOX.⁴⁵⁻⁵² Curcumin shows hepato-protective, cyto-protective, neuro-protective and antioxidant properties.^{44,49,50,53-56}



Turmeric (*Curcuma longa*)

Green Tea (*Camellia sinensis*)

With a 5,000-year history as valued beverage and medicine, tea prepared from *Camellia sinensis* is one of the most widely consumed beverages in the world. Polyphenols comprise 30% to 40% of green tea leaves. These are flavonols and flavanols (catechins) – including epigallocatechin-3-gallate (EGCG), which impart significant antioxidant and anti-inflammatory activity.⁵⁷⁻⁶⁰ Tea polyphenols benefit cell cycle regulation and are found to be cell-protective.⁶¹⁻⁶³ EGCG, the main catechin in green tea extract, inhibits VEGF (vascular endothelial growth factor)^{64,65} implicated in abnormal cell behavior, cell signaling and angiogenesis.⁶⁵⁻⁶⁸

Japanese Knotweed (*Polygonum cuspidatum*)

Polygonum has been used widely for centuries in Chinese medicine to prevent and treat disease. In modern research it is found to exert antimicrobial, anti-inflammatory, neuro-protective and cardio-protective functions. Knotweed contains quercetin, catechin and resveratrol, which all demonstrate powerful antioxidant and anti-inflammatory properties. Knotweed extract is found to inhibit inflammatory pathways including TNF- α , NF- κ B and COX-2 expression.^{69,70}



Knotweed (*Polygonum cuspidatum*)

Resveratrol

An active constituent of Knotweed, resveratrol, is extensively researched for its profound benefits. Numerous studies report its role in preventing age-related disorders including diabetes, cancer, cardiovascular and neuro-degenerative conditions.^{43,71} Research reports that resveratrol binds to numerous cell-signaling molecules, activates transcription factors, induces antioxidant enzymes, modulates cell-cycle regulation and inhibits expression of inflammatory processes.^{72,73} Due to its inherent properties as a polyphenolic compound, resveratrol is neuro-protective and can help inhibit progression of age-related neurological decline and disease. A potent antioxidant, resveratrol works through multiple cellular pathways to support normal function.⁷⁴ Studies show that resveratrol indirectly inhibits NF- κ B signaling.²

Amla (*Emblica officinalis*)

Amla fruit, also known as Indian Gooseberry, is a valued Ayurvedic restorative and rejuvenative tonic. Used for thousands of years as food and medicine, it supplies a rich source of vitamin C, flavonoids and carotenoids.⁷⁵ Amla fruit is traditionally given to relieve circulatory, digestive and respiratory conditions. Studies find it possesses potent antioxidant and anti-inflammatory activity.⁷⁶⁻⁷⁹ Amla has an ORAC (oxygen radical absorbance capacity) value of 1770 – twice that of Acai berry and about 17 times that of pomegranate fruit.⁷⁹ Amla fruit contains gallic acid, ellagic acid and sesquiterpenoids, which are known to be free-radical scavenging, antioxidant, anti-inflammatory, antimutagenic and immunomodulatory.⁸⁰⁻⁸²

Fatty Acids: Dietary Foundation

Fatty acid intake (quality and quantity) is well-known to exert major influence on function and modulation of healthy inflammatory, immune and hormonal processes. Fatty acids can drive either pro- or anti-inflammatory processes and influence the AA cascade. EPA (eicosapentaenoic acid) and DHA (docosahexaenoic acid) – metabolites of omega-3 fatty acids – compete with the omega-6 arachidonic acid for conversion by COX enzymes into prostaglandins (PGs). Omega-3 oil metabolism results in higher production of anti-inflammatory PGs, while that of omega-6 results in formation of pro-inflammatory PGs.

An optimal dietary ratio of around 2:1 of omega-6 to -3 fatty acids is found to help suppress inflammation and provide a wide array of health benefits.⁸³ In the Western world, the dietary ratio is often about 15:1 – reflecting over-consumption of omega-6 oils along with pronounced deficiency of omega-3 oils.⁸³ This is widely recognized as a major contributory factor to diseases including cardiovascular, cellular dysfunction, autoimmune and inflammatory conditions.⁸³⁻⁸⁵ Historically, fish constitutes the major dietary source of omega-3 oils. Important plant sources include flax, chia and hemp seeds and walnuts. Supplementing a healthy diet with fish and plant oils further enhances a balanced intake of the much-needed omega-3 oils.

Fish Oils

Fish oils from small fish, including krill, anchovy and sardines, provide essential fatty acids including EPA and DHA. Omega-3 fish oils inhibit COX-2 expression along with the oxidative metabolism of AA to the inflammatory PGs.^{86,87} Omega-3 fatty acids can benefit serum lipid levels and help reduce C-reactive protein levels.⁸⁸ EPA is found to improve the prognosis of chronic inflammatory diseases including atherosclerosis.⁸⁹⁻⁹¹

Studies find that EPA and DHA benefit bone mineral density in those with rheumatoid arthritis.⁹¹ Fish oils benefit those with joint diseases such as arthritis^{92,93} and can help reduce neck and back pain, likely through their influence in calming the inflammatory process.⁹⁴

Sea Buckthorn Oil (*Hippophae rhamnoides*)

Sea Buckthorn oil (SBO) contains a unique fatty acid profile, offering a nutritious blend of omega-3, -6, -7 and -9 unsaturated fatty acids. It is one of only a few plants that is especially rich in the omega-7 fatty acid palmitoleic acid. The omega-7 group of fatty acids is of interest to researchers for its ability to calm inflammation and support multiple physiological functions. The potent antioxidant activity attributed to Sea Buckthorn derives partially from its highly concentrated content of flavonoids, vitamin C, tocopherols, tocotrienols and a rich array of carotenoids.⁹⁵⁻⁹⁷



Sea Buckthorn (*Hippophae rhamnoides*)

SBO enhances healthy glutathione levels at the cellular level along helps reduce production of reactive oxygen species.⁹⁸ In many studies, SBO was found to significantly influence the glutathione redox system, increasing available and circulating levels of glutathione.⁹⁹⁻¹⁰⁰

Herbs and Natural Compounds to Calm Inflammation and Pain

Natural foods, spices and herbs high in polyphenols provide the foundation of a healthy lifestyle and diet. A formulation of polyphenolic-rich botanicals provides a potent multi-targeted approach to calm inflammation and enhance restorative physiological functions. The addition of adaptogenic formulas enhances restoration of healthy allostasis and benefits any therapeutic approach. Focused botanical formulas chosen for their specific anti-inflammatory influence contributes a multi-faceted and synergistic approach to address chronic inflammation.

Pain, often a significant component of the chronic inflammatory process, is usually associated with injury or disease. Chronic pain affects around 100 million Americans – a greater number than those affected by diabetes, heart disease and cancer combined.¹⁰¹ The most common chronic pain conditions include migraines and severe headaches, back pain, neck pain, facial pain, cancer pain, joint pain and neurogenic pain.¹⁰²

Chronic pain involves non-resolving inflammation and continual firing of pain signals in the nervous system. Typically, the sensation of pain is transmitted by afferent neurons from the periphery of the body to the spinal cord and from there to the brain, eliciting various responses, including production of natural endorphins. Feedback loops modulate and modify this process. Inflammatory processes and

neurotransmitters are implicated in the complex pain response. Recent studies report that spinal glial-mediated neuro-inflammation is a key factor in the development of chronic pain. This involves production of chemokines via NF-kB which plays a major role in the inflammatory response.^{18,19,103}

In current pain research, botanicals are being widely studied in the search for compounds that naturally exert a multi-targeted approach. Scientists often discover the advantage of using whole herbs, as each herb contains a complex array of biochemical compounds that naturally form a synergistic and multi-targeted effect. These herbs work on pathways other than the opioid receptors, for example through modulating dopamine and cannabinoid receptors^{104,105} or by mediating neuroglial inflammation, particularly in the spine.^{18,19}

This section discusses potent traditional herbs that are highly researched and recognized for their ability to work through multiple pathways to exert potent anti-inflammatory influence and help to alleviate pain.

Frankincense (*Boswellia serrata*)

Frankincense contains gum-like resinous constituents known as boswellic acids, which exhibit powerful anti-inflammatory, analgesic and health-promoting activity. Boswellic acids are the key compounds in Frankincense that are most studied, including one known as AKBA (acetyl-11-keto-beta-boswellic acid).

Boswellia extract is found to benefit those with inflammatory bowel conditions.¹⁰⁶ Boswellic acid extracts are found to down-regulate multiple inflammatory pathways including 5-LOX, COX-2 and NF-kB. LOX acts as a biological fuel for cellular dysfunction by stimulating EGF (epidermal growth factor), VEGF (vascular endothelial growth factor) and other growth factors.^{36,107,108} These extracts are also found to induce apoptosis, modulate cell-signaling^{36,109,110} and to exert immunomodulatory influence.¹¹¹⁻¹¹³

Feverfew (*Tanacetum parthenium*)

This member of the Aster family is traditionally used to prevent migraine headaches.¹¹⁴ Ancient Greeks and Europeans used it to reduce fever and to calm inflammation in disorders including psoriasis, rheumatism and colic. Feverfew is high in flavonoids and sesquiterpene lactones; both of which are known for their anti-inflammatory qualities. Parthenolide, one of the sesquiterpene lactones in Feverfew, is of great interest to researchers as it is found to interfere with the inflammatory actions of AA, histamine and NF-kB.^{115,116} Parthenolide inhibits the 5-LOX and COX pathways and helps prevent conversion of AA to prostaglandins.¹¹⁷ It inhibits pro-inflammatory, cytokine-mediated cell-signaling and inhibits PG synthesis through various pathways. It is thought to benefit migraines through multiple pathways including through inhibition of PG synthesis and by decreasing spasm of vascular smooth muscles.¹¹⁸



Feverfew (*Tanacetum parthenium*)

Magnolia (*Magnolia officinalis*)

Magnolia bark, widely used in Chinese medicine, is rich in biologically-active compounds including alkaloids, coumarins, flavonoids, lignans and terpenoids.¹¹⁹ Its main active constituent is considered to be the polyphenol honokiol, which exhibits potent antioxidative and anti-inflammatory

activity. Research finds that Magnolia bark extract strongly inhibits COX-2 pathways, formation of prostaglandins, TNF, NF-kB and IL (interleukin) factors.¹²⁰⁻¹²³ Honokiol is being studied for its neuro-protective influence as it is found to calm oxidative and inflammatory processes in neurons and microglial cells.¹¹⁹

Andrographis (*Andrographis paniculata*)

Both Chinese and Ayurvedic medicines highly regard this herb for its ability to alleviate inflammatory and infectious diseases. Andrographis is high in flavonoids and diterpenoids. This herb acts as antipyretic and anti-inflammatory and is able to modulate both the humoral and cellular adaptive immune response. Andrographis is found to inhibit NO (nitric oxide) and PG production¹²⁴⁻¹²⁸ along with expression of COX-2.^{126,129} Andrographis extract is noted for its ability to reduce lipid peroxidation and inhibit formation of reactive oxygen species. It also enhances the antioxidant enzyme SOD (superoxide dismutase).^{128,130}



Andrographis (*Andrographis paniculata*)

Ginger (*Zingiber officinalis*)

Ginger is known and used worldwide as cooking spice, herbal home remedy and revered botanical medicine. It is a daily household remedy for digestive upset, sore throat, colds and flu. Ginger aids circulation and has a thermogenic and diaphoretic effect. It shows significant antioxidant¹³¹⁻¹³³ and anti-inflammatory activity. Ginger is found to inhibit COX-2 expression and NF-kB activation.¹³⁴⁻¹³⁵ Ginger also influences PG metabolism, inhibits thromboxane synthesis and is found to significantly inhibit platelet aggregation and inflammation.¹³⁶⁻¹³⁸

Long Pepper (*Piper longum*) and Piperine from Black Pepper (*Piper nigrum*)

Long Pepper is one of the world's oldest culinary and medicinal spices. It was much more widely known, used and traded than Black Pepper until around the 14th century. Studies report that Long Pepper is antioxidant, anti-inflammatory and immunomodulatory.^{40,139-143}

Black Pepper berries naturally contain the alkaloid piperine, a powerful and highly-researched compound. Piperine is known to be antioxidative, antimutagenic, antibacterial and hepatoprotective.^{144,145} Studies find piperine helps reduce levels of pro-inflammatory mediators including COX-2, IL factors and TNF-alpha. It also supports healthy glutathione and SOD levels.^{146,147} It is found to inhibit VEGF and to modulate cytokine and growth factor responses.¹⁴³

Corydalis Root (*Corydalis spp.*)

Corydalis is the premier herb in traditional Chinese medicine for pain of all kinds especially when combined with appropriate herbs. Modern studies report that Corydalis demonstrates potent analgesic and anti-inflammatory activity. Two alkaloids isolated from Corydalis, THP (tetrahydropalmatine) and DHCB (dehydrocorybulbine), demonstrate significant analgesic and sedative effects on the central nervous system.^{102,148-150} In one study, 75mg THP showed nerve pain reduction in 78% of patients tested.¹⁵⁰

THP and DHCB are found to exert a mild agonist effect on the m opioid receptor. The affinity of DHCB to dopamine receptors is found to be greater than 100 times its affinity for the m opioid receptor.¹⁰² THP and DHCB, structurally similar, bind to dopamine receptors where they are potent

antagonists. Many studies show dopamine plays a role in modulating pain circuits in the brain. Studies with mice engineered to be deficient in dopamine were found to be more sensitive to pain suggesting that dopamine is involved with pain threshold.¹⁰⁴

The alkaloid DHCB exerts analgesic effects independent of the opioid receptors and is found effective in both inflammatory and neuropathic pain models with repeated use. The Herbalome project is currently working on a DHCB-based drug for pain.^{102,151} Researchers also find Corydalis root used as a whole herb extract is highly efficacious.¹⁵² The analgesic effect of Corydalis involves participation of the cannabinoid receptors.¹⁵³

White Willow Bark (*Salix alba*)

White Willow bark, used for thousands of years by many cultures, is known to alleviate pain, calm inflammation and reduce fever. Hippocrates (400 BC) was said to advise his patients to chew on the bark to reduce fever and inflammation. In modern times, German chemist Felix Hoffman first isolated the compound salicin from the bark. This was later modified to the chemical acetylsalicylic acid, or aspirin. Salicin, a potent anti-inflammatory agent, inhibits over-expression of COX-2 and NF- κ B factors which are implicated in inflammation and abnormal gene expression.^{154,155} The advantage of White Willow bark extract is that it does not irritate the stomach lining. This is because the salicin naturally found in White Willow bark is only converted to the acid form after absorption by the stomach.



White Willow Bark (*Salix alba*)

Native Americans used White Willow bark to calm joint pain. Modern studies confirm its effectiveness and safety as an analgesic giving relief in arthritic joint pain, low back pain, headaches and low back pain.¹⁵⁶⁻¹⁶⁰

Notoginseng (*Panax notoginseng*)

Chinese medicine highly reveres Notoginseng as the primary herb for all traumatic injuries including falls, fractures, contusions and pains. It is used both internally and externally as a hemostatic after traumatic injuries. Modern research find that Notoginseng is rich in saponins and immune-enhancing polysaccharides.¹⁶¹ It is found to have the ability to address hypoxia and anoxia.¹⁶² As a potent anti-inflammatory, Notoginseng modulates various inflammatory pathways and exerts pain-relieving effect in animal studies.¹⁶³⁻¹⁶⁶

White Peony Root (*Paeonia lactiflora*)

In Chinese medicine, Peony is a major tonic herb. Peony is found to contain flavonoids, proanthocyanidins, tannins, terpenoids, complex polysaccharides and glycosides, of which paeoniflorin is the most studied. Paeoniflorin, monoterpenoids and other constituents of White Peony are known to be spasmolytic.¹⁶⁷⁻¹⁶⁹ The compound paeoniflorin (PF) is found to calm pain and joint swelling and to lessen the decline of bone and cartilage in experimental arthritis. It suppresses multiple inflammatory processes including prostaglandin and leukotriene production, and production of reactive oxygen species, proinflammatory cytokines and chemokines. It inhibits formation of new blood vessels and production of matrix metalloproteinases. It is found beneficial to relieve both signs and symptoms of rheumatoid arthritis.¹⁷⁰

PF is a monoterpene glycoside that is also neuro-protective. It is found to calm the microglial inflammatory response, to reduce production of pro-inflammatory factors from activated microglial cells and to calm inflammation-induced neurotoxicity.^{171,172} Another constituent of White Peony, albiflorin, is found to exert anti-inflammatory benefits similar to paeoniflorin.¹⁷³

Dong Quai Root (*Angelica sinensis*)

One of the most esteemed herbs in the Chinese materia medica, Dong Quai is especially revered as a nourishing tonic. Studies find that the analgesic and anti-inflammatory effects of Dong Quai are similar to those of acetylsalicylic acid. A preparation of Dong Quai showed 82.9% effective rate in 35 patients with migraine headache.¹⁷⁴

High in coumarins and flavonoids, Dong Quai also contains volatile oils, vitamin A, carotenoids, vitamins B12, E and phytosterols (including beta-sitosterol) along with minerals. The root does not exert estrogenic activity.¹⁷⁵ Dong Quai is known for hematopoietic, antioxidant, immune regulatory properties and is neuro-protective.¹⁷⁶ A major component of Dong Quai, known as ligustilide (LIG), is shown to have neuro-protective and anti-inflammatory effects. Studies show that spinal astrocyte-mediated neuro-inflammation plays an important role in the pathogenesis of chronic pain. The spinal cord mediates the pain response and it is found that calming chemokine production in the spinal nerves helps decrease overall pain.^{18,19,177} LIG exerts an anti-inflammatory influence, inhibiting NF-κB-mediated chemokine production in spinal astrocytes, and demonstrates effectiveness for inflammatory pain.¹⁷⁸

Conclusion: Botanicals and Natural Compounds Modulate Inflammatory Response

Foods, spices and botanical medicines have been used safely and effectively for thousands of years to enhance health, calm inflammation and promote the healing response. Many botanicals are able to calm the inflammatory response through their dynamic interaction with the epigenetic and immune systems. Found to work through multiple pathways, they influence diverse cellular pathways to restore allostasis.

The epigenetic, immune and inflammatory responses are but a few aspects of this ongoing process of allostasis. Because of its inherent plasticity and far-reaching regulatory effects, epigenetic modulation is currently considered a highly-effective therapeutic target. It is theorized that botanicals offer a diverse complexity of molecular compounds that interface harmoniously with human physiology and molecular dynamics. There are no single or linear responses in human physiology; rather a continual, dynamic of cellular interactions, cascades and signaling. Botanical medicines demonstrate the ability to exert a modulatory effect and help promote resolution of chronic inflammation and restoration of allostasis.

Offering a broad spectrum of natural compounds with natural affinity for human physiology, botanical medicines are well-tolerated. Combination therapies are found to be beneficial either through adjunctive or synergistic enhancement. As researchers investigate the numerous compounds found in botanicals, they often discover that use of the actual plant material, and a combination of botanicals, offers comprehensive and synergistic therapeutic benefit.^{2,9,10,12,24,31,37,39,40,152}

For more information, please call 1-888-628-8720 or email info@naturahhealthproducts.com.



References

1. McEwen BS. *Protective and damaging effects of stress mediators: central role of the brain*. Dialogues in Clinical Neuroscience 2006. Vol 8(4):367-381.
2. Salminen A, Kaarniranta K. *Genetics vs. entropy: Longevity factors suppress the NF- κ B-driven entropic aging process*. Ageing Research Reviews 2010. (9)298–314.
3. Wierda RJ, Geutskens SB, et al. *Epigenetics in atherosclerosis and inflammation*. J Cell Mol Med. Epigenetics Review Series 2010. Vol 14(6A):1225-1240.
4. Yasmin R, Siraj S, et al. *Epigenetic regulation of inflammatory cytokines and associated genes in human malignancies*. Mediators of Inflammation 2015. Hindawi Publishing Corp. Article ID 201703. 8 pages. <http://dx.doi.org/10.1155/2015/201703>
5. Martel-Pelletier J, Lajeunesse D et al. *Therapeutic role of dual inhibitors of 5-LOX and COX, selective and non-selective non-steroidal anti-inflammatory drugs*. Ann Rheum Dis 2003. 762: 501-509.
6. Bayarsaihan D. *Epigenetic mechanisms in inflammation*. Critical Reviews in Oral Biology & Medicine. J Dent Res 2011. 90(1):9-17.
7. Shanmugam MK, Sethi G. *Role of epigenetics in inflammation-associated diseases*. Subcell Biochem 2013. 61:627-57. doi: 10.1007/978-94-007-4525-4_27.
8. Shanmugam MK, Sethi G. Chapter 27: *Role of Epigenetics in Inflammation-Associated Diseases*. in: Epigenetics: Development and Disease. Volume 61 of the series Subcellular Biochemistry.
9. Pandey K, Rizvi S. *Plant polyphenols as dietary antioxidants in human health and disease*. Oxidative Medicine and Cellular Longevity. November/December 2(5) 270-278.
10. Bharat B, Aggarwal RV et al. *Targeting Inflammatory Pathways for Prevention and Therapy of Cancer: Short-Term Friend, Long-Term Foe*. Clin Cancer Res 2009. 15(2).
11. Ricciotti E, FitzGerald GA. *Prostaglandins and Inflammation*. Arterioscler Thromb Vasc Biol. 2011 May. 31(5): 986-1000. doi:10.1161/ATVBAHA.110.207449.
12. Lawrence T. *The Nuclear Factor NF- κ B Pathway in Inflammation*. Cold Spring Harb Perspect Biol 2009. 1:a001651
13. Copstead LC, Banasik JL. *Pathophysiology*, Fifth Edition. 2013. Elsevier Saunders.
14. Alberts B, Johnson A et al. *Molecular Biology of the Cell*, 4th Edition. 2002. Garland Science.
15. Dubois RN, Abramson SB, Crofford L, et al. *Cyclooxygenase in biology and disease*. The FASEB J. Sept 1998. Vol.12: 1063-1073.
16. Solana R, Mariani E. *NK and NK/T cells in human senescence*. Vaccine. 2000 Feb 25. 18(16):1613-20.
17. Ji RR, Xu ZZ and Gao YJ. *Emerging targets in neuroinflammation-driven chronic pain*. Nature Reviews Drug Discovery. 2014. (13) 533–548. doi:10.1038/nrd4334. Published online 20 June 2014.
18. Zhu MD, Zhao LX, et al. *Ligustilide inhibits microglia-mediated proinflammatory cytokines production and inflammatory pain*. Brain Res Bull. 2014 Oct. 109:54-60. doi: 10.1016/j.brainresbull.2014.10.002. Epub 2014 Oct 16.
19. Zhao LX, Jiang BC, et al. *Ligustilide attenuates inflammatory pain via inhibition of NF- κ B-mediated chemokines production in spinal astrocytes*. Eur J Neurosci. 2014 Apr. 39(8):1391-402. doi: 10.1111/ejn.12502. Epub 2014 Feb 12.
20. Duque GA, Descoteaux A. *Macrophage cytokines: involvement in immunity and infectious diseases*. Frontiers in Immunology. 07 Oct 2014. doi: 10.3389/fimmu.2014.00491
21. Stow JL, Low PC, et al. *Cytokine secretion in macrophages and other cells: pathways and mediators*. Immunobiology 2009. 214(7):601-12. doi: 10.1016/j.imbio.2008.11.005. Epub 2009 Mar 5.
22. Stow JL, Murray RZ. *Intracellular trafficking and secretion of inflammatory cytokines*. Cytokine Growth Factor Rev. 2013 Jun. 24(3):227-39. doi: 10.1016/j.cytogfr.2013.04.001. Epub 2013 May 4.
23. Kronfol Z, Remick DG. *Cytokines and the brain: implications for clinical psychiatry*. Am J Psychiatry 2000. 157:683-694.
24. Spelman K, Burns JJ, et al. *Modulation of cytokine expression by traditional medicines: a review of herbal immunomodulators*. Alt Med Rev 2006. Vol 11(2) 128-150.
25. Venkatraman M, Anto RJ, Nair A, et al. *Biological and chemical inhibitors of NF- κ B sensitized SiHa cells to cisplatin-induced apoptosis*. Mol Carcinog. 2005 Jul 25. 44(1):51-59.

26. Fujisawa S, Atsumi T, et al. *Cytotoxicity, ROS-generation activity and radical-scavenging activity of curcumin and related compounds*. *Anticancer Res*. 2004 Mar-Apr. 24(2B):563-9.
27. Nakshatri H, Goulet RJ. *NF-kappaB and breast cancer*. *Curr Probl Cancer*. 2002 Sep-Oct. 26(5):282-309. Review.
28. Samuelsson B. *Arachidonic acid metabolism: role in inflammation*. *Z Rheumatol* 1991. 50 Suppl 1:3-6.
29. Khanapure SP, Garvey DS, et al. *Eicosanoids in inflammation: biosynthesis, pharmacology, and therapeutic frontiers*. *Curr Top Med Chem*. 2007. 7(3):311-40.
30. Clària J. *Cyclooxygenase-2 biology*. *Curr Pharm Des* 2003. 9(27):2177-90.
31. Joseph J et al. *Reversing the deleterious effects of aging on neuronal communication and behavior: beneficial properties of fruit polyphenolic compounds*. *Am J Clin Nutr* 2005. 81(suppl):313S-6S.
32. Yao X, Li H, Leng SX. *Inflammation and Immune System Alterations in Frailty*. *Clin Geriatr Med*. 2011 Feb. 27(1): 79-87. doi:10.1016/j.cger.2010.08.002
33. Li H, Manwani B, Leng SX. *Frailty, inflammation, and immunity*. *Aging Dis*. 2011 Dec. 2(6):466-73. Epub 2011 Dec 2.
34. Müller M, Kersten S. *Nutrigenomics: Goals and Perspectives*. *Nature Reviews Genetics*. 2003. (4):315 -322.
35. Orzechowski A, Ostaszewski P, et al. *Bioactive substances of plant origin in food--impact on genomics*. *Reprod Nutr Dev*. 2002 Sep-Oct. 42(5):461-77. Review. Erratum in: *Reprod Nutr Dev*. 2002 Nov-Dec. 42(6):625.
36. Wallace JM. *Nutritional and botanical modulation of the inflammatory cascade--eicosanoids, cyclooxygenases, and lipoxygenases--as an adjunct in cancer therapy*. *Integr Cancer Ther*. 2002 Mar. 1(1):7-37.
37. Anand P, Sundaram C, et al. *Curcumin and cancer: an "old-age" disease with an "age old" solution*. *Cancer Letters* 2008. 267:133-164.
38. Manach, C, Scalberri, A, Morand, C, et al. *Polyphenols: food sources and bioavailability*. *Am J Clin Nutr*. May 2004. Vol 79:5: 727-747.
39. Tsao R. *Chemistry and Biochemistry of Dietary Polyphenols*. *Nutrients* 2010. (2):1231-1246. doi:10.3390/nu2121231
40. Akbar P, Jahan I, et al. *Antioxidant capacity of piper longum and piper nigrum fruits grown in Bangladesh*. *World Journal of Pharmaceutical Sciences* ISSN (Print): 2321-3310; ISSN (Online): 2321-3086. Atom and Cell Publishers. Available online at: <http://www.wjpsonline.org/>
41. Ahmad N, Fazal H, et al. *Biological role of Piper nigrum L. (Black pepper): A review*. *Asian Pac J of Trop Biomedicine* 2012. S1945-S1953.

Turmeric

42. Sikha A, Harini A, Hegde Prakash L. *Pharmacological activities of wild turmeric (Curcuma aromatica Salisb): a review*. *Journal of Pharmacognosy and Phytochemistry* 2015. 3(5): 01-04.

43. Kunnumakkara AB, Diagaradjane P, et al. *Curcumin sensitizes human colorectal cancer xenografts in nude mice to gamma-radiation by targeting nuclear factor-kappaB-regulated gene products*. *Clin Cancer Res*. 2008 Apr 1. 4(7):2128-36.

44. Katori M, Jajima M. *Multiple roles of inducible cyclooxygenase-2 and its selective inhibitors*. *Nippon Yakurigaku Zasshi*.1997. 109(6):247:58.

45. *Inhibition of tumor necrosis factor by curcumin, a phytochemical*. *Biochemical Pharmacology (United Kingdom)*. 1995. 49/11:1551-1556.

46. Funk JL, Oyarzo JN, et al. *Turmeric extracts containing curcuminoids prevent experimental rheumatoid arthritis*. *J Nat Prod*. 2006 Mar. 69(3):351-5.

47. Van't Land B, Blijlevens NM, et al. *Role of curcumin and the inhibition of NF-kappaB in the onset of chemotherapy-induced mucosal barrier injury*. *Leukemia*. 2004 Feb. 18(2): 276 84.

48. Hussain AR, Ahmed M, et al. *Curcumin suppresses constitutive activation of nuclear factor-kappa B and requires functional Bax to induce apoptosis in Burkitt's lymphoma cell lines*. *Mol Cancer Ther*. 2008 Oct 7. (10):3318-29.

49. Bagchi D, Preuss, HG. *Phytopharmaceuticals in Cancer Chemoprevention. Chapter 23, Curcumin Derived from Turmeric (Curcuma longa): a Spice for all Seasons*. 2005 CRC Press, Boca Raton, FL. p349-387.

50. Jiang MC, Yang-Yen HF, et al. *Curcumin induces apoptosis in immortalized NIH 3T3 and malignant cancer cell lines*. *Nutr Cancer* 1996. 26:111-20.

51. Venkatraman M, Anto RJ, Nair A, et al. *Biological and chemical inhibitors of NF-kappaB sensitize SiHa cells to cisplatin-induced apoptosis*. *Mol Carcinog*. 2005 Jul 25. 44(1):51-59.

52. Fujisawa S, Atsumi T, et al. *Cytotoxicity, ROS-generation activity and radical-scavenging activity of curcumin and related compounds*. *Anticancer Res*. 2004 Mar-Apr. 24(2B):563-9.



53. *Anti-tumour and antioxidant activity of natural curcuminoids*. Cancer Lett (IRELAND) Jul 20 1995. 94 (1):79-83.

54. Zbarsky V, Datla KP, Parkar S, et al. *Neuroprotective properties of the natural phenolic antioxidants curcumin and naringenin but not quercetin and fisetin in a 6-OHDA model of Parkinson's disease*. Free Radic Res. 2005 Oct. 39(10):1119-25.

55. Mythri RB, Jagatha B, et al. *Mitochondrial Complex I Inhibition in Parkinson's Disease: How Can Curcumin Protect Mitochondria?* Antioxid Redox Signal. 2007 Jan.

56. Mythri RB, Bharath MM. *Curcumin: a potential neuroprotective agent in Parkinson's disease*. Curr Pharm Des 2012;18(1):91-9.

Green Tea

57. Sarkar A, Bhaduri A. *Black Tea Is a Powerful Chemopreventor of Reactive Oxygen and Nitrogen Species: Comparison with Its Individual Catechin Constituents and Green Tea*, Biochemical and Biophysical Research Communications. June 1, 2001. Vol. 284, No. 1.

58. Higashi-Okai K et al. *Identification and antioxidant activity of several pigments from the residual green tea (Camellia sinensis) after hot water extraction*, J UOEH, 2001Dec 1. 23(4):335-44.

59. Henning SM, Fajardo-Lira C, et al. *Catechin content of 18 teas and a green tea extract supplement correlates with the antioxidant capacity*. Nutr Cancer 2003. 45(2):226-35.

60. Blumberg J. *Tea flavonoids gain recognition as powerful weapon against disease*. J Nutrition. 1 October 2003. Vol 133, No 10.

61. Kakuda T. *Neuroprotective effects of the green tea components theanine and catechins*. Biol Pharm Bull 2002. 25:1513-1518.

62. Ahn WS, Yoo J, Huh SW, et al. *Protective effects of green tea extracts (polyphenon E and EGCG) on human cervical lesions*. Eur J Cancer Prev. 2003 Oct. 12(5):383-90.

63. Ahmad N, et al. *Green tea constituent epigallocatechin-3 galate and induction of apoptosis and cell cycle arrest in human carcinoma cells*. J National Cancer Inst 1997. 89(24):1881-1886.

64. Sartippour MR, et al. *Green Tea Inhibits Vascular Endothelial Growth Factor (VEGF) Induction in Human Breast Cancer Cells*, Department of Surgery, Division of Oncology and Center for Human Nutrition, University of California, Los Angeles, CA and University of Texas, MD Anderson Cancer Center, Houston, TX.

65. Sartippour MR, Shao ZM, et al. *Green tea inhibits vascular*

endothelial growth factor (VEGF) induction in human breast cancer cells. J Nutr 2002. 132(8):2307-11.

66. Lee SR, Im KJ, Suh SI, Jung JG. *Protective effect of green tea polyphenol (-)-epigallocatechin gallate and other antioxidants on lipid peroxidation in gerbil brain homogenates*. Phytother Res 2003 Mar. 17(3):206.

67. Rodriguez SK, Guo W, Liu L, et al. *Green tea catechin, epigallocatechin-3-gallate, inhibits vascular endothelial growth factor angiogenic signaling by disrupting the formation of a receptor complex*. Int J Cancer. 2006 Apr 1. 118(7):1635-44.

68. Shankar S, Ganapathy S, Srivastava RK. *Green tea polyphenols: biology and therapeutic implications in cancer*. Front Biosci. 2007 Sep 1. 12:4881-4899.

Knotweed

69. Zhang H, Li C, et al. *A Review of the Pharmacological Effects of the Dried Root of Polygonum cuspidatum (Hu Zhang) and Its Constituents*. Evid Based Complement Alt Med. 2013. 2013:208349. doi: 10.1155/2013/208349. Epub 2013 Sep 30.

70. Feng L, Zhang LF, et al. *Studies on active substance of anticancer effect in Polygonum cuspidatum*. Zhong Yao Cai. 2006 Jul. 29(7):689-91.

Resveratrol

71. Turner, RS, Thomas, RD, et al. *A randomized, double-blind, placebo-controlled trial of resveratrol for Alzheimer disease*. Neurology. pub online Sep 11, 2015 DOI 10.1212/WNL.0000000000002035

72. Kulkarni SS, Cantó C. *The molecular targets of resveratrol*. Biochim Biophys Acta. 2015 Jun. 1852(6):1114-23. doi: 10.1016/j.bbadis.2014.10.005. Epub 2014 Oct 12.

73. Harikumar KB, Aggarwal BB. *Resveratrol: a multitargeted agent for age-associated chronic diseases*. Cell Cycle. 2008 Apr 15. 7(8):1020-35. Epub 2008 Feb 15.

74. Zhuang H, Kim YS, et al. *Potential mechanism by which resveratrol, a red wine constituent, protects neurons*. Ann N Y Acad Sci. 2003 May. 993:276-86: discussion 287-8.

Amla

75. Scartezzini P, Antognoni F, et al. *Vitamin C content and antioxidant activity of the fruit and of the Ayurvedic preparation of Emblica officinalis Gaertn*. J Ethnopharmacol. 2006 Mar 8. 104(1-2):113-8. Epub 2005 Oct 13.

76. Takagi N, Sanashiro T. *Health foods containing antioxidative and anti-allergy food materials*, Jpn. Kokai Tokkyo Koho 1996. JP 10 00,070



77. Yokozawa T, Kim HY, Kim HJ, et al. *Amla (Emblca officinalis Gaertn.) prevents dyslipidaemia and oxidative stress in the ageing process*. Br J Nutr. 2007 Jun. 97(6):1187-95.

78. Naik GH, Priyadarsini KI, Bhagirathi RG, et al. *In vitro antioxidant studies and free radical reactions of triphala, an ayurvedic formulation and its constituents*. Phytother Res. 2005 Jul. 19(7):582-6.

79. McNally, A. *Indian gooseberry - the new superfruit?* 2001/2008 – Decision News Media SAS 29-Jan-2008.

80. Baliga MS, Dsouza JJ (2011). *Amla (Emblca officinalis Gaertn), a wonder berry in the treatment and prevention of cancer*. Eur J Cancer Prev. 2011 May. 20(3):225-39. doi: 10.1097/CEJ.0b013e32834473f4.

81. Chen TS, Liou SY, Wu HC, et al. *Efficacy of Epigallocatechin-3-Gallate and Amla (Emblca officinalis) Extract for the Treatment of Diabetic-Uremic Patients*. J Med Food. 2011 Jul-Aug. 14(7-8):718-23. Epub 2011 Jun 1.

82. Huang JL, Zhong ZG. *Study of galic acid extracted from the leaves of Phyllanthus emblica on apoptotic mechanism of human hepatocellular carcinoma cells BEL-7404*. Zhong Yao Cai. 2011 Feb. 34(2):246-9. Chinese.

Fish Oils

83. Simopoulos AP. *The importance of the ratio of omega-6/omega-3 essential fatty acids*. Biomed Pharmacother. 2002 Oct. 56(8):365-79.

84. Watkins B, Seifert M. *Omega-3s crucial to bone health in later years*. J Nutrit Biochem. 13/07/2005. Purdue University's Center for Enhancing Foods to Protect Health.

85. Yamagishi K, Iso H, et al. JACC Study Group. *Fish, omega-3 Polyunsaturated Fatty Acids, and Mortality From Cardiovascular Diseases in a Nationwide Community-Based Cohort of Japanese Men and Women* The JACC (Japan Collaborative Cohort Study for Evaluation of Cancer Risk) Study. J Am Coll Cardiol. 2008 Sep 16. 52(12):988-996.

86. McCarty MF. *Fish oil may impede tumour angiogenesis and invasiveness by down-regulating protein kinase C and modulating eicosanoid production*. Med. Hypotheses 1996. 46: 107-115.

87. Calder PC. *Dietary modification of inflammation with lipids*. Proc Nutr Soc 2002: 61:345-58.

88. Ambring A, Johansson M, et al. *Mediterranean-inspired diet lowers the ratio of serum phospholipid n-6 to n-3 fatty acids, the number of leukocytes and platelets, and vascular endothelial growth factor in healthy subjects*. Am J Clin Nutr. 2006 Mar. 83(3):575-81.

89. Matsuyama W, Mitsuyama H, et al. *Effects of omega-3 polyunsaturated fatty acids on inflammatory markers in COPD*. Chest. 2005 Dec.128(6):3817-27.

90. Thies F, Garry JM, et al. *Association of n-3 polyunsaturated fatty acids with stability of atherosclerotic plaques: a randomised controlled trial*. Lancet. 2003 Feb 8. 361(9356):477-85.

91. Erkkila, A. *Fish intake is associated with a reduced progression of coronary artery atherosclerosis in postmenopausal women with coronary artery disease*. Am J Clin Nutr 2004. 80:626-32.

92. Bhattacharya A, Rahman M, et al. *Inhibition of osteoporosis in autoimmune disease prone MRL/Mpj-Fas(lpr) mice by N-3 fatty acids*. J Am Coll Nutr. 2005 Jun. 24(3):200-9.

93. Proudman SM, James MJ, et al. *Fish oil in recent onset rheumatoid arthritis: a randomised, double-blind controlled trial within algorithm-based drug use*. Ann Rheum Dis. 2013 Sept 30.

94. Maroon JC, Bost JW. *Omega-3 fatty acids (fish oil) as an anti-inflammatory: an alternative to nonsteroidal anti-inflammatory drugs for discogenic pain*. Surg Neurol. 2006 Apr. 65(4):326-31.

Sea Buckthorn Oil

95. Bernstein, Adam M. et al. *Purified palmitoleic acid for the reduction of high-sensitivity C-reactive protein and serum lipids: A double-blinded, randomized, placebo controlled study*. J Clin Lipidology. Vol 8(6):612-617.

96. Yang B, Kalimo KO, et al. *Effects of dietary supplementation with sea buckthorn (Hippophae rhamnoides) seed and pulp oils on atopic dermatitis*. J Nutr Biochem. 1999 Nov. 10(11):622-30.

97. Larmo P, Alin J, Salminen E, et al. *Effects of sea buckthorn berries on infections and inflammation: a double-blind, randomized, placebo-controlled trial*. European J Clin Nutr. Vol. 86: 2345-2353. Advance online publication 27 June 2007. doi: 10.1038/sj.ejcn.1602831

98. Upadhyay NK, Kumar R, et al. *Safety and healing efficacy of Sea buckthorn (Hippophae rhamnoides L.) seed oil on burn wounds in rats*. Food Chem Toxicol. 2009 Jun. 47(6):1146-53.

99. Wu D, Meng Z. *Effect of sulfur dioxide inhalation on the glutathione redox system in mice and protective role of sea buckthorn seed oil*. Arch Environ Contam Toxicol. 2003 Oct. 45(3):423-8.

100. Ruan A, Min H, Meng Z, Lu Z. *Protective effects of seabuckthorn seed oil on mouse injury induced by sulfur dioxide inhalation*. Inhal Toxicol. 2003 Sep. 15(10):1053-8.



Pain

101. American Academy of Pain Medicine: *AAPM Facts and Figures on Pain*. http://www.painmed.org/patientcenter/facts_on_pain.aspx

102. Zhang Y, Wang C, et al. *A Novel Analgesic Isolated from a Traditional Chinese Medicine*. *Current Biol*. Jan 20, 2014. (24):117-123. Elsevier. <http://dx.doi.org/10.1016/j.cub.2013.11.039>

103. Ji RR, Xu ZZ and Gao YJ. *Emerging targets in neuroinflammation-driven chronic pain*. *Nature Reviews Drug Discovery* 2014. (13) 533-548. doi:10.1038/nrd4334. Pub online 20 June 2014.

104. Ingram SL. *Pain: Identification of novel analgesics from traditional Chinese medicines*. *Curr Biol*. 2014 February 3. 24(3): R114-R116. doi:10.1016/j.cub.2013.12.030.

105. Huang JY, Fang M, et al. *Analgesic effect of Corydalis yanhusuo in a rat model of trigeminal neuropathic pain*. *Nan Fang Yi Ke Da Xue Xue Bao*. 2010 Sep. 30(9):2161-4.

Boswellia

106. Gupta I, Parihar A, Malhotra P, et al. *Effects of gum resin of Boswellia serrata in patients with chronic colitis*. *Planta Med*. 2001 Jul. 67(5):391-5.

107. Safayhi H, Sailer ER, Ammon HP. *Mechanism of 5-lipoxygenase inhibition by acetyl-11-keto-beta-boswellic acid*. *Mol Pharmacol*. 1995 Jun. 47(6):1212-6.

108. Takada Y, Ichikawa H, et al. *Acetyl-11-keto-beta-boswellic acid potentiates apoptosis, inhibits invasion, and abolishes osteoclastogenesis by suppressing NF-kappa B and NF-kappa B-regulated gene expression*. *J Immunol*. 2006 Mar 1. 176(5):3127-40.

109. Park YS, Lee JH, Harwalkar JA, et al. *Acetyl-11-keto-beta-boswellic acid (AKBA) is cytotoxic for meningioma cells and inhibits phosphorylation of the extracellular-signal regulated kinase 1 and 2*. *Adv Exp Med Biol*. 2002. 507:387-93.

110. Yuan Y, Cui SX, Wang Y, et al. *Acetyl-11-keto-beta-boswellic acid (AKBA) prevents human colonic adenocarcinoma growth through modulation of multiple signaling pathways*. *Biochim Biophys Acta*. 2013 Oct. 1830(10):4907-16. doi:10.1016/j.bbagen.2013.06.039. Epub 2013 Jul 10.

111. Badria FA, Mikhaeil BR, et al. *Immunomodulatory triterpenoids from the oleogum resin of Boswellia carterii* Birdwood. *Z Naturforsch*. 2003 Jul-Aug. 58(7-8):505-16.

112. Mikhaeil BR, Maatooq GT, et al. *Chemistry and immunomodulatory activity of frankincense oil*. *Z Naturforsch*.

2003 Mar-Apr. 58(3-4):230-8.

113. Ammon HP. *Modulation of the immune system by Boswellia serrata extracts and boswellic acids*. *Phytomedicine*. 2010 Sep. 17(11):862-7. Epub 2010 Aug 8.

Feverfew

114. Prusinski A, Durko A, Niczyporuk-Turek A. *Feverfew as prophylactic treatment of migraine*. *Neurol Neurochir Pol* 1999. 33 Suppl 5:89-95.

115. Pareek A., Suthar M et al. *Feverfew (Tanacetum parthenium L.): A systematic review*. *Pharmacogn Rev*. 2011 Jan-Jun; 5(9):103-110. doi: 10.4103/0973-7847.79105.

116. Cutlan AR, Bonilla LE, et al. *Intra-specific variability of feverfew: correlations between parthenolide, morphological traits and seen origin*. *Planta Med* 2000 Oct; 66(7):612-7.

117. Zhang S, Lin ZN, Yang CF, et al. *Suppressed NF-kappaB and sustained JNK activation contribute to the sensitization effect of parthenolide to TNF-alpha-induced apoptosis in human cancer cells*. *Carcinogenesis*. 2004 Nov. 25(11):2191-9. Epub 2004 Jul 15.

118. Yip KH, Zheng MH, Feng HT, et al. *Sesquiterpene lactone parthenolide blocks lipopolysaccharide-induced osteolysis through the suppression of NF-kappaB activity*. *J Bone Miner Res*. 2004 Nov. 19(11):1905-16.

Magnolia

119. Chuang DY, Chan MH, Zong Y et al. *Magnolia polyphenols attenuate oxidative and inflammatory responses in neurons and microglial cells*. *J Neuroinf* 2013. 10:15.

120. Kim BH, Cho JY. *Anti-inflammatory effect of honokiol is mediated by PI3K/Akt pathway suppression*. *Acta Pharmacol Sin*. 2008 Jan. 29(1):113-22.

121. Lee E, Jung J. et al. *Anti-inflammatory effects of magnolol and honokiol are mediated through inhibition of the downstream pathway of MEKK-1 in NF-kappaB activation signaling*. *Planta Med* 2005. 71:338-343.

122. Tse AK, Wan CK, Shen XL, et al. *Honokiol inhibits TNF-alpha-stimulated NF-kappaB activation and NF-kappaB-regulated gene expression through suppression of IKK activation*. *Biochem Pharmacol*. 2005 Nov 15; 70(10):1443-57. Epub 2005 Sep 21.

123. Ahn KS, Sethi G, Shishodia S, et al. *Honokiol potentiates apoptosis, suppresses osteoclastogenesis, and inhibits invasion through modulation of nuclear factor-kappaB activation pathway*. *Mol Cancer Res*. 2006 Sep. 4(9):621-33.

Andrographis

124. Mills S and Bone K. *Principles and Practices of Phytotherapy*. London, New York; Churchill Livingstone, 2001.

125. Gonzalez PA et al. *Andrographolide interferes with T cell activation and reduces experimental autoimmune encephalomyelitis in the mouse*. J Pharmacol Exp Ther. 2005;312. 1:366-72.

126. Liu J, Wang ZT, et al. *Inhibitory effects of neoandrographolide on nitric oxide and prostaglandin E2 production in LPS-stimulated murine macrophage*. Mol Cell Biochem. 2007 Apr. 298(1-2):49-57. Epub 2006 Nov 16.

127. Liu J, Wang ZT, Ji LL. *In vivo and in vitro anti-inflammatory activities of neoandrographolide*. Am J Chin Med 2007. 35(2):317-28.

128. Sheeja K, Shihab PK, Kuttan G. *Antioxidant and anti-inflammatory activities of the plant Andrographis paniculata* Nees. Immunopharmacol Immunotoxicol 2006. 28(1):129-40.

129. Shen T, Yang WS, Yi YS, et al. *AP-1/IRF-3 Targeted Anti-Inflammatory Activity of Andrographolide Isolated from Andrographis paniculata*. Evid Based Comp Alt Med 2013. 210736. doi: 10.1155/2013/210736. Epub 2013 Jun 6.

130. Akbar, S. *Andrographis paniculata: A review of pharmacological activities and clinical effects*. Alt Med Rev. 2011. Vol 16(1): 66-77.

Ginger

131. Guo P, Xu J, et al. *Inhibition of hydrogen peroxide production in chondrocytes induced by fulvic acid by ginger volatile oil*. China J Chinese Materia Medica 1997. 22:559-61.

132. Zhou Y, Xu R. *Antioxidative effect of Chinese drugs*. Chung Kuo Chung Yao Tsa Chih 1992: 17:368-9, 373.

133. Cao ZF, Chen ZG, Guo P, et al. *Scavenging effects of ginger on superoxide anion and hydroxyl radical*. Zhongguo Zhong Yao Za Zhi. 1993 Dec. 18(12):750-1, 764.

134. Kim SO, Chun KS, et al. *Inhibitory effects of [6]-gingerol on PMA-induced COX-2 expression and activation of NF-kappaB and p38 MAPK in mouse skin*. Biofactors. 2004. 21(1-4):27-31.

135. Kim SO, Kundu JK, Shin et al. *Gingerol inhibits COX-2 expression by blocking the activation of p38 MAP kinase and NF-kappaB in phorbol ester-stimulated mouse skin*. Oncogene. 2005 Apr 7. 24(15):2558-67.

136. Kiuchi F, Shibuya M, Sankawa U. *Inhibitors of*

prostaglandin biosynthesis from ginger. Chem Pharm Bull. Tokyo 1982. 30:754-7.

137. Flynn D, Rafferty M, Boctor A. *Inhibition of human neutrophil 5-lipoxygenase activity by gingerdione, shogaol, capsaicin and related pungent compounds*. Prostaglandins Leukotrienes Med 1986. 24:195-8.

138. Srivastava KC. *Isolation and effects of some ginger components of platelet aggregation and eicosanoid biosynthesis*. Prostaglandins. Leukot Med 1986. 25:187-98.

Long Pepper

139. Manoj P, Soniya EV et al. *Recent studies on well-known spice, Piper longum Linn*. Natural product Radiance. July-August 2004. Vol 3(4):222-.

140. Kumar S, Kamboj J et al. *Overview for Various Aspects of the Health Benefits of Piper Longum Linn. Fruit*. J Acupunct Meridian Stud 2011. 4(2):134-140.

141. Zaveril M, Khandhar A, et al. *Chemistry and pharmacology of Piper longum L*. Int J Pharm Sci Rev and Res. Nov/Dec 2010. Vol 5, Issue 1. Article-010.

142. Vasavirama K, Upender M. *Piperine: A valuable alkaloid from Piper species*. Int J Pharm Pharm Sci. Vol 6(4): 34-38.

143. Sunila ES, Kuttan G. *Piper longum inhibits VEGF and proinflammatory cytokines and tumor-induced angiogenesis in C57BL/6 mice*. Int Immunopharmacology 6:2006:733– 741.

Black Pepper/Piperine

144. Ahmad N, Fazal H et al. *Biological role of Piper nigrum L. (Black pepper): A review*. Asian Pacific Journal of Tropical Biomedicine. 2012. S1945-S1953.

145. Matsuda H, Ninomiya K, et al. *Hepatoprotective amide constituents from the fruit of Piper chaba: Structural requirements, mode of action, and new amides*. Bioorg Med Chem. 2009 Oct 15. 17(20):7313-23. Epub 2009 Aug 29.

146. Umar S, Golam Sarwar AH, Umar K, et al. *Piperine ameliorates oxidative stress, inflammation and histological outcome in collagen induced arthritis*. Cell Immunol. 2013 Jul 19. 284(1-2):51-59. doi: 10.1016/j.cellimm.2013.07.004.

147. Ying X, Chen X, Cheng S, et al. *Piperine inhibits IL-β induced expression of inflammatory mediators in human osteoarthritis chondrocyte*. Int Immunopharmacol. 2013 Oct. 17(2):293-9. doi: 10.1016/j.intimp.2013.06.025. Epub 2013 Jul 6.

Corydalis

148. Hsu HY. *Oriental Materia Medica: A Concise Guide*. Long Beach, CA. Oriental Healing Arts Institute 1986. p448–50.



149. Zhu YP. *Chinese Materia Medica: Chemistry, Pharmacology, and Applications*. Australia. Harwood Academic Publishers 1998. p445–8.

150. Lin MT, Wang JJ, Young MS. *The protective effect of dl-tetrahydropalmatine against the development of amygdala kindling seizures in rats*. *Neurosci Lett* 2002 Mar 8. 320(3):113-6.

151. Zhang X, Liu Y, et al. *The herbalome—an attempt to globalize Chinese herbal medicine*. *Analytical and Bioanalytical Chemistry*. January 2012. Vol 402(2):573.

152. University of California Irvine. *Chinese herbal compound relieves inflammatory and neuropathic pain*. Online article. January 2014. <http://news.uci.edu/press-releases/chinese-herbal-compound-relieves-inflammatory-and-neuropathic-pain/>

153. Huang JY, Fang M, et al. *Analgesic effect of Corydalis yanhusuo in a rat model of trigeminal neuropathic pain*. *Nan Fang Yi Ke Da Xue Xue Bao*. 2010 Sep. 30(9):2161-4.

White Willow Bark

154. Chrubasik S, Pollak S, Black A. *Willow bark extract, a useful alternative for the treatment of osteoarthritis: comment on the editorial by Marcus and Suarez-Almazor*. *Arthritis Rheum*. 2003 Jan. 48(1):278-80.

155. Bonaterra GA, Heinrich EU, et al. *Anti-inflammatory effects of the willow bark extract STW 33-I (Proaktiv((R))) in LPS-activated human monocytes and differentiated macrophages*. *Phytomedicine* May 28, 2010.

156. Schmid B, Ludtke R, Selbmann HK, et al. *Effectiveness and tolerance of standardized willow bark extract in arthrosis patients, Randomized, placebo controlled double-blind study*. *Z Rheumatol*. 2000 Oct. 59(5):314-20.

157. Chrubasik S, Kunzel O, Model A, et al. *Treatment of low back pain with a herbal or synthetic anti-rheumatic: a randomized controlled study*. *Willow bark for low back pain*. *Rheumatology* 2001. 40:1388-1393.

158. Chrubasik S, Eisenberg E, Balan E, et al. *Treatment of low back pain exacerbations with willow bark extract: a randomized double-blind study*. *Am J Med*. 2000 Jul. 109(1):9-14.

159. Gagnier JJ, van Tulder M, et al. *Herbal medicine for low back pain*. *Cochrane Database Syst Rev*. 2006 Apr 19(2):CD004504.

160. Uehleke B, Müller J, Stange R, et al. *Willow bark extract STW 33-I in the long-term treatment of outpatients*

with rheumatic pain mainly osteoarthritis or back pain. *Phytomedicine*. 2013 Aug 15. 20(11):980-4. doi: 10.1016/j.phymed.2013.03.023. Epub 2013 Jun 2.

Notoginseng

161. Gao H, Wang F, et al. *Immunostimulating polysaccharides from Panax notoginseng*. *Pharmaceutical Research* 1996. 13(8):1196-1200.

162. Cui XM, Lo CK, et al. *Authentication of Panax notoginseng by 5S-rRNA spacer domain and random amplified polymorphic DNA (RAPD) analysis*. *Planta Medica* 2003. 69(6): 584-586.

163. Wang YL, Chen D, Wu JL. *Effects and mechanism of total saponins of Panax Notoginseng on anti-inflammation and analgesia*. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 1994. 14(1): 35-36.

164. Chang SH, Choi Y, Park JA, et al. *Anti-inflammatory effects of BT-201, an n-butanol extract of Panax notoginseng, observed in vitro and in a collagen-induced arthritis model*. *Clinical Nutrition* 2007. 26(6): 785-791.

165. Zhang JH, Wang JP, Wang HJ. *Clinical study on effect of total panax notoginseng saponins on immune related inner environment imbalance in rheumatoid arthritis patients*. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 2007. 27(7): 589-592.

166. Jung HW, Seo UK, et al. *Flower extract of Panax notoginseng attenuates lipopolysaccharide-induced inflammatory response via blocking of NF-kappaB signaling pathway in murine macrophages*. *Journal of Ethnopharmacology* 2009. 122(2): 313-319.

Peony

167. *Monograph: Peony (Paeonia spp.)*. *Alt Med Rev*. 2001. Vol 6 (5):495-499

168. Kimura M, Kimura I, Takahashi K, et al. *Blocking effects of blended paeoniflorin or its related compounds with glycyrrhizin on neuromuscular junctions in frogs and mice*. *Japan J Pharmacol* 1984. 36:275-282.


169. Zhang XJ, Chen HL, Li Z. *Analgesic effect of paeoniflorin in rats with neonatal maternal separation-induced visceral hyperalgesia is mediated through adenosine A(1) receptor by inhibiting the extracellular signal-regulated protein kinase (ERK) pathway*. *Pharmacol Biochem Behav*. 2009 Nov. 94(1):88-97. doi: 10.1016/j.pbb.2009.07.013. Epub 2009 Aug 5.

170. Zhang W, Dai SM. *Mechanisms involved in the therapeutic effects of Paeonia lactiflora Pallas in rheumatoid arthritis*. *Int Immunopharmacol*. 2012 Sep. 14(1):27-31. doi: 10.1016/j.intimp.2012.06.001. Epub 2012 Jun 21.



171. Nam KN, Yaw CG, Hong JW, et al. *Paeoniflorin, a monoterpene glycoside, attenuates lipopolysaccharide-induced neuronal injury and brain microglial inflammatory response*. Biotechnol Lett. 2013 Aug. 35(8):1183-9. doi: 10.1007/s10529-013-1192-8. Epub 2013 Apr 5.
172. He DY, Dai SM. *Anti-inflammatory and immunomodulatory effects of paeonia lactiflora pall., a traditional chinese herbal medicine*. Front Pharmacol. 2011 Feb 25. 2:10. doi: 10.3389/fphar.2011.00010. eCollection 2011.
- 173 55. Wang QS, Gao T, et al. *Comparative studies of paeoniflorin and albiflorin from Paeonia lactiflora on anti-inflammatory activities*. Pharm Biol. 2014 Sep. 52(9):1189-95. doi: 10.3109/13880209.2014.880490. Epub 2014 Mar 19.
- Dong Quai**
174. Chen JK, Chen TT. *Chinese Medical Herbology and Pharmacology*. Art of Medicine Press. 2004.
175. *Monograph. Angelica sinensis (Dong quai)*. Alt Med Rev. 2004. Volume 9, Number 4.
176. Chen XP, Li W, et al. *Phytochemical and pharmacological studies on Radix Angelica sinensis*. Chin J Nat Med. 2013 Nov. 11(6):577-87. doi: 10.1016/S1875-5364(13)60067-9.
177. Xu J, E XQ, et al. *Angelica Sinensis attenuates inflammatory reaction in experimental rat models having spinal cord injury*. Int J Clin Exp Pathol. 2015 Jun 1. 8(6):6779-85. eCollection 2015.
178. Qian B, Li F, et al. *Ligustilide Ameliorates Inflammatory Pain and Inhibits TLR4 Upregulation in Spinal Astrocytes Following Complete Freund's Adjuvant Peripheral Injection*. Cell Mol Neurobiol. 2015 Jun 27.

About the Authors



Donald R. Yance Jr., RH (AHG), CN

Donald R. Yance Jr., RH (AHG), CN, is a practicing Clinical Master Herbalist, Certified Nutritionist and the author of two books including his latest *Adaptogens in Medical Herbalism* (Healing Arts Press, 2013). Providing visionary leadership in the field of natural medicine, Yance has maintained a successful clinical practice for over two decades and trains practitioners worldwide in his therapeutic methodology for patient care. He is the founder and formulator of Natura Health Products, a line of botanical and nutritional formulas, which are based on key principles of Yance's Eclectic Triphasic Medical System.



Suzanne E. Sky, L.Ac., MTOM

Suzanne E. Sky, L.Ac., MTOM has been a Chinese medicine practitioner since 1989 and involved with herbs, nutrition and the healing arts since the mid-70s. She worked with Donald Yance as clinical associate for five years until 2004 and since then often as a writing associate.

With Chinese medicine as her framework, Suzanne integrates herbal, nutritional and lifestyle recommendations along with gentle energy work (Jin Shin Jyutsu) or acupuncture. Blending ancient Chinese wisdom with modern knowledge, her work focuses on supporting and nourishing each person's innate healing and regenerative capacity. Her clinic, Ashland Acupuncture, is located in Ashland, Oregon. She teaches Qi Gong and Chinese medicine classes.